

Geology

The leading edge of the Greater Himalayan Crystalline complex revealed in the NW Indian Himalaya: Implications for the evolution of the Himalayan orogen

A. Alexander G. Webb, An Yin, T. Mark Harrison, Julien C  l  rier and W. Paul Burgess

Geology 2007;35:955-958

doi: 10.1130/G23931A.1

Email alerting services

click www.gsapubs.org/cgi/alerts to receive free e-mail alerts when new articles cite this article

Subscribe

click www.gsapubs.org/subscriptions/ to subscribe to *Geology*

Permission request

click <http://www.geosociety.org/pubs/copyrt.htm#gsa> to contact GSA

Copyright not claimed on content prepared wholly by U.S. government employees within scope of their employment. Individual scientists are hereby granted permission, without fees or further requests to GSA, to use a single figure, a single table, and/or a brief paragraph of text in subsequent works and to make unlimited copies of items in GSA's journals for noncommercial use in classrooms to further education and science. This file may not be posted to any Web site, but authors may post the abstracts only of their articles on their own or their organization's Web site providing the posting includes a reference to the article's full citation. GSA provides this and other forums for the presentation of diverse opinions and positions by scientists worldwide, regardless of their race, citizenship, gender, religion, or political viewpoint. Opinions presented in this publication do not reflect official positions of the Society.

Notes

The leading edge of the Greater Himalayan Crystalline complex revealed in the NW Indian Himalaya: Implications for the evolution of the Himalayan orogen

A. Alexander G. Webb*

An Yin

T. Mark Harrison

Julien C el erier

W. Paul Burgess

 Department of Earth and Space Sciences and Institute of Geophysics and Planetary Physics,
 University of California–Los Angeles, Los Angeles, California 90095-1567, USA

Research School of Earth Sciences, Australia National University, Canberra, ACT 2601, Australia

 Department of Earth and Space Sciences and Institute of Geophysics and Planetary Physics,
 University of California–Los Angeles, Los Angeles, California 90095-1567, USA

ABSTRACT

The three Himalayan lithologic units, the Lesser Himalayan Sequence, the Greater Himalayan Crystalline complex, and the Tethyan Himalayan Sequence, have a specific structural correlation with the Main Central thrust and South Tibet detachment in the central Himalaya. There, the Main Central thrust places the Greater Himalayan Crystalline complex over the Lesser Himalayan Sequence, and the South Tibet detachment places the Tethyan Himalayan Sequence over the Greater Himalayan Crystallines. Although this division has formed the basis for all Himalayan tectonic models, it fails to explain aspects of the geology of the western Himalaya where the Main Central thrust places the Tethyan Himalayan Sequence directly above the Lesser Himalayan Sequence. Our mapping in NW India shows that this relationship results from southward merging of the Main Central thrust and South Tibet detachment. This finding, in conjunction with observed alternating shear senses on the South Tibet detachment, is inconsistent with the wedge-extrusion and erosion-induced channel-flow models (both require only top-to-the-N motion on the South Tibet detachment) but is consistent with a tectonic-wedging model.

Keywords: Himalaya, South Tibet detachment, Main Central thrust, tectonic wedge.

INTRODUCTION

The ~2500-km-long Himalayan orogen is widely thought to consist of only three major units: the Lesser Himalayan Sequence (LHS; mainly low-grade Proterozoic metasediments), the Greater Himalayan Crystalline complex (GHC; largely high-grade paragneisses and migmatite), and the Tethyan Himalayan Sequence (THS; dominantly low-grade late Proterozoic to Eocene shelf sediments) (Heim and Gansser, 1939). In the central Himalaya, the LHS and GHC are separated by the Main Central thrust, and the GHC and THS are separated by the South Tibet detachment (LeFort, 1996) (Fig. 1). However, in the western Himalaya (west of 77°E), the Main Central thrust places THS rocks directly over LHS metasediments (e.g., Yeats and Lawrence, 1984; Frank et al., 1995; Pogue et al., 1999) (Fig. 1). Several scenarios have been advanced to explain this different relationship (e.g., Thakur, 1998; DiPietro and Pogue, 2004; Yin, 2006), but uncertainty regarding the position of the South Tibet detachment in many locations in the NW Indian Himalaya (cf. Fig. 1 of Searle et al. [1999] and plate 1 of Steck [2003]) limits efforts to understand its significance. This paper summarizes the results of field work undertaken in the western Himalaya that lead to an interpretation that explains this relationship but that challenges current views of Himalayan thrust tectonics.

GEOLOGY OF THE ROHTANG LA AREA

Our field area is located in the Rohtang La area northwest of the Kulu Window (Figs. 1 and 2). Although the South Tibet detachment can be traced from Nepal to this area, its westward extension is poorly defined (Fig. 2) (Choudhuri et al., 1992; Vannay and Grasemann, 1998; Jain et al., 1999), which had led to various interpretations including connection with the Zaskar shear zone (Searle et al., 1999) and termination in the Rohtang La area (Steck, 2003). We mapped the position of the South Tibet detachment by tracking its deformation zone, metamorphic grade changes across the fault, and marker beds along the fault.

Typically, the South Tibet detachment shear zone is hundreds of meters thick and exhibits both top-to-the-NE and top-to-the-SW shear-sense indicators (also see Jain et al., 1999). This contrasts to the top-to-the-SW motion associated with the Main Central thrust ductile shear zone below. In general, gneisses are common below and schists are prevalent above the South Tibet detachment. Although garnet is present both above and below the South Tibet detachment, kyanite and/or sillimanite are diagnostic of the South Tibet detachment footwall. Following Vannay and Steck (1995) and Wyss et al. (1999), we used graphitic quartzite and discontinuous lenses of calc-silicate schists in the THS as marker beds to trace the South Tibet detachment hanging wall. Intrusive contacts around Cambrian-Ordovician granites in the South Tibet detachment hanging wall are undeformed, whereas the same contacts in the South Tibet detachment footwall are intensely transposed by ductile folding.

At Rohtang La, the South Tibet detachment shear zone preserves ductile shear fabrics including top-to-the-SW S-C fabric, top-to-the-NE sigma augen, top-to-the-NE and top-to-the-SW shear band cleavage, and top-to-the-SW folds (Fig. 3). The top-to-the-NE shear fabrics overprint top-to-the-SW shear fabrics (also see Jain et al., 1999). A sharp contact between mylonitic augen gneiss below and garnet schist above is present in the South Tibet detachment shear zone, which we interpret as the South




Figure 1. Map of central and western Himalaya compiled from DiPietro and Pogue (2004), Valdiya (1980), Yin (2006), and references for Figure 2. GCT—Great Counter Thrust; GHC—Greater Himalayan Crystalline complex; LHS—Lesser Himalayan Sequence; MCT—Main Central thrust; Q—Quaternary alluvium; SH—Sub-Himalayan Sequence; STD—South Tibet detachment; THS—Tethyan Himalayan Sequence; ZSZ—Zaskar shear zone.

*E-mail: awebb@ess.ucla.edu.




Figure 2. Map of Kulu Window region (after Frank et al., 1973, 1975; Sharma, 1977; Vannay and Steck, 1995; Vannay and Grasemann, 1998; Wyss et al., 1999; our own observations, and additional references in the GSA Data Repository¹).

Tibet detachment fault. The South Tibet detachment can be traced south of Rohtang La into the south-verging overturned Phojal anticline (i.e., the Phojal Nappe of Frank et al., 1973) along the west bank of the Beas River (Fig. 3A). The South Tibet detachment is overturned in the Phojal anticline (Figs. 3A and 3B) as indicated by the folding of (1) the gneiss-schist contact that marks the fault and features top-to-the-NE sigma augen and top-to-the-NE C' shear band cleavage (Fig. 3C), (2) the hanging-wall marker units of graphitic quartzite and calc-silicate schist, and (3) mineral isograds directly above and below the fault (e.g., Frank et al., 1973; Epard et al., 1995). Although the original geometry of the isograds may not have been subhorizontal (e.g., Le Fort, 1975), the subparallel relationship between our mapped South Tibet detachment and the regional isograds north of the anticline (i.e., Frank et al., 1973) is consistent with overturned folding of the South Tibet detachment at this location.

The overturned South Tibet detachment can be traced southeastward from the Phojal anticline to the northern edge of the Main Central thrust Kulu Window near Manikaran (Fig. 3A). Using our lithologic and metamorphic criteria, we have extended the mapped location of the South Tibet detachment southeast from Manikaran for another 50 km by correlation with the same gneiss-schist contact mapped by Sharma (1977). Still farther east, the overturned South Tibet detachment must merge with the Main Central thrust along the northeastern margin of the Kulu Window because the schist in the South Tibet

detachment hanging wall pinches out west of the well-studied Sutlej River section where Greater Himalayan Crystalline gneiss is exposed continuously between the Main Central thrust and the South Tibet detachment (e.g., Vannay and Grasemann, 1998) (Fig. 2). Thus, the merging of the South Tibet detachment and Main Central thrust in map view defines the tip line of a southward-tapering GHC wedge (Fig. 3B). The overturned South Tibet detachment and Phojal anticline are eroded over the Kulu Window and reappear at the southeast side of the window, which we mapped along the Pabbar River (Fig. 2). The overturned South Tibet detachment there can be well constrained by metamorphic-grade variation across the fault and the South Tibet detachment hanging-wall marker units.

DISCUSSION

The South Tibet detachment makes a sharp U-turn in map view at Rohtang La, changing from a gently west-dipping structure to a NE-dipping overturned fault that merges with the Main Central thrust (Fig. 2). Overturned folding may have been caused by distributed top-to-the-SW shear across the Main Central thrust zone, suggesting that the South Tibet detachment became inactive prior to cessation of motion on the Main Central thrust. This is consistent with age constraints showing that both faults were active in the early Miocene, but Main Central thrust slip may extend to the middle Miocene (Catlos et al., 2002; Vannay et al., 2004).

The three-layer division of the Himalaya has been explained by wedge-extrusion, channel-flow, and general-shear models (Burchfiel and Royden, 1985; Grujic et al., 1996; Vannay and Grasemann, 2001; Nelson et al., 1996; Beaumont et al., 2001). Note that the Nelson et al. (1996) channel-flow model differs from that of

Beaumont et al. (2001) in that the latter predicts India crust to be subducted and then return to the surface without transporting Asian rocks to the present Himalayan range. This physical process may be more appropriately characterized as corner flow (see Cloos, 1982). None of these models explain our observed southward merging of the Main Central thrust and South Tibet detachment, but instead require the Main Central thrust and South Tibet detachment to be surface faults. The tunneling stage of channel flow of Beaumont et al. (2001) is compatible with the observed Main Central thrust–South Tibet detachment branch line geometry, but fails to explain two key kinematic observations. First, its predicted top-to-the-N South Tibet detachment kinematics is inconsistent with the observed alternating top-to-the-N and top-to-the-S shear fabrics in the South Tibet detachment zone in NW India and throughout the Himalaya (e.g., Patel et al., 1993; Hodges et al., 1996; Grujic et al., 2002; Robinson et al., 2006; this study). Second, channel-flow tunneling requires slip on the South Tibet detachment and Main Central thrust to vanish at their branch line where the tunnel terminates, which is inconsistent with >100 km of Main Central thrust slip south of the Main Central thrust–South Tibet detachment branch line as constrained by the distance between the Kulu Window and the Simla Klippe (Fig. 1).

To reconcile the new observation for the Main Central thrust–South Tibet detachment relationship and the alternating shear motion on the South Tibet detachment, we propose that the GHC was emplaced as a tectonic wedge (see Fig. 10 of Price, 1986) via southward slip on the Main Central thrust and alternating top-to-the-N and top-to-the-S motion on the South Tibet detachment (Fig. 4) (Yin, 2006). Depending on the displacement boundary condition at the back side of the GHC thrust sheet, sense of shear along the South Tibet detachment could have alternated between top-to-the-N and top-to-the-S (Figs. 4B and 4C). Even when motion along the South Tibet detachment was top-to-the-N, the THS moved southward with respect to the LHS (Figs. 4D, 4E, and 4F). The top-to-the-N South Tibet detachment motion may link to top-to-the-N slip on the Great Counter thrust along the Indus-Tsangpo suture (Figs. 1 and 4H) (e.g., Yin et al., 1994). The Great Counter thrust forms the roof thrust of a second south-directed tectonic wedge, the Asia Plate (see Fig. 4H). The alternating insertion of the two tectonic wedges could have produced the temporally varying shear sense on the South Tibet detachment (Figs. 4A–C and 4H).

Although some parts of the Himalaya show multiple alternations in shear-sense along the South Tibet detachment (e.g., Hodges et al., 1996), the dominant pattern is a sequential change from top-to-the-S to top-to-the-N shear on the South Tibet detachment (e.g., Patel et al.,

¹GSA Data Repository item 2007235, additional references in support of Figure 2, is available online at www.geosociety.org/pubs/ft2007.htm, or on request from editing@geosociety.org or Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.




Figure 3. A: Geologic map of Rohtang La area, based on our mapping, as highlighted by strike and dip symbols, and a compilation of work from Frank et al. (1973, 1995), Vannay and Steck (1995), Wyss et al. (1999), and Wyss (2000). B: Geologic cross section of Rohtang La area. C: Top-to-the-NNE mylonitic gneiss, view to WNW. Location is shown in A.

1993; Jain et al., 1999; Grujic et al., 2002). Uniformly top-to-the-N South Tibet detachment slip could produce this pattern because records of top-to-the-S shear from the upper part of the wedge-front shear zone (Main Central thrust-south) would have been transported to the wedge-top shear zone (South Tibet detachment) across the Main Central thrust-South Tibet detachment branch line (see Figs. 4D–4F). Thus, the early Main Central thrust top-to-the-S shear fabrics would have been overprinted by the later South Tibet detachment top-to-the-N shear fabrics.

The discovery of the Main Central thrust–South Tibet detachment branch line has important implications for explaining along-strike variation of the Himalayan geology and its relationship to exhumation. Because the erosional pattern of the Himalaya may be asymmetric, with an eastward increase in the magnitude of exhumation (Finlayson et al., 2002), it is possible that the Main Central thrust–South Tibet detachment branch line is preserved in the western Himalaya but eroded away in the central Himalaya (Fig. 1).




Figure 4. Tectonic wedge model for GHC emplacement, involving top-to-the-S Main Central thrust slip and alternating top-to-the-N and top-to-the-S slip along STD. A: Predeformation geometry. B: GHC emplacement during top-to-the-S faulting along South Tibet detachment. C: GHC emplacement during top-to-the-N faulting along South Tibet detachment. Despite top-to-the-N relative motion along South Tibet detachment, note that THS is consistently thrust south with respect to LHS. D–F: Top-to-the-N slip along South Tibet detachment transfers records of top-to-the-S shear from Main Central thrust hanging wall to South Tibet detachment hanging wall. Records of top-to-the-S shear from Main Central thrust-south are shown in gray. In inset, active shear indicators are shown in black. G: Late distributed shear during motion on Main Central thrust may overturn the South Tibet detachment. H: Schematic diagram of early Miocene development of Himalaya, involving two tectonic wedges (Price, 1986) inserted to south: GHC and Asia plate. Thrust emplacement of both wedges can produce temporally varying shear sense along South Tibet detachment. GCT—Great Counter Thrust; GHC—Greater Himalayan Crystalline complex; LHS—Lesser Himalayan Sequence; MCT—Main Central thrust; STD—South Tibet detachment; THS—Tethyan Himalayan Sequence.

CONCLUSIONS

Field mapping in the NW India Himalaya reveals southward-up merging of the South Tibet detachment and Main Central thrust and a complex South Tibet detachment slip history alternating between top-to-the-N and top-to-the-S shear. These observations are inconsistent with existing Himalayan models, but they are consistent with a tectonic-wedging model (Price, 1986). They also help to explain the change in structural correlation between the major Himalayan faults (Main Central thrust and South Tibet detachment) and the Himalayan units (Lesser Himalayan Sequence, Greater Himalayan Crystalline complex, and Tethyan Himalayan Sequence) from the central to western Himalaya: the Main Central thrust places the GHC over the LHS in the central Himalaya, but it juxtaposes the THS over the LHS in the western Himalaya. This variation can be attributed to an eastward increase in the magnitude of exhumation resulting in differential preservation of past orogenic architecture. This implies that the along-strike variation of Himalayan geology may not be a result of a change in deformation mechanism but a consequence of spatially varying erosion.

ACKNOWLEDGMENTS

This research was supported by the National Science Foundation (NSF) Tectonics Program. Webb was supported by an NSF Graduate Research Fellowship. We thank Peter DeCelles, Joseph DiPietro, Mike Murphy, Lothar Ratschbacher, and an anonymous reviewer for excellent and constructive reviews.

REFERENCES CITED

- Beaumont, C., Jamieson, R.A., Nguyen, M.H., and Lee, B., 2001, Himalayan tectonics explained by extrusion of a low-viscosity crustal channel coupled to focused surface denudation: *Nature*, v. 414, p. 738–742, doi: 10.1038/414738a.
- Burchfiel, B.C., and Royden, L.H., 1985, North-south extension within the convergent Himalayan region: *Geology*, v. 13, p. 679–682, doi: 10.1130/0091-7613(1985)13<679:NEWTCH>2.0.CO;2.
- Catlos, E.J., Harrison, T.M., Manning, C.E., Grove, M., Rai, S.M., Hubbard, M.S., and Upreti, B.N., 2002, Records of the evolution of the Himalayan orogen from in situ Th-Pb ion microprobe dating of monazite: Eastern Nepal and western Garhwal: *Journal of Asian Earth Sciences*, v. 20, p. 459–479, doi: 10.1016/S1367-9120(01)00039-6.
- Choudhuri, B.K., Bist, K.S., and Rawat, B.S., 1992, Evidence for thrusting between the Higher Himalayan Crystallines and Tethyan metasediments in Lahaul and Kullu Valleys, Himachal Pradesh: *Journal of Himalayan Geology*, v. 3, p. 191–194.
- Cloos, M., 1982, Flow mélanges: Numerical modeling and geologic constraints on their origin in the Franciscan subduction complex, California: *Geological Society of America Bulletin*, v. 93, p. 330–345.
- DiPietro, J.A., and Pogue, K.R., 2004, Tectonostratigraphic subdivisions of the Himalaya: A view from the west: *Tectonics*, v. 23, p. TC5001, doi: 10.1029/2003TC001554.
- Epard, J.L., Steck, A., Vannay, J.C., and Hunziker, J., 1995, Tertiary Himalayan structures and metamorphism in the Kulu Valley (Mandi-Khoksar transect of the Western Himalaya)—Shikar-Beh-Nappe and Crystalline Nappe: *Schweizerische Mineralogische und Petrographische Mitteilungen*, v. 75, p. 59–84.
- Finlayson, D.P., Montgomery, D.R., and Hallet, B., 2002, Spatial coincidence of rapid inferred erosion with young metamorphic massifs in the Himalayas: *Geology*, v. 30, p. 219–222, doi: 10.1130/0091-7613(2002)030<0219:SCORIE>2.0.CO;2.
- Frank, W., Hoinkes, G., Miller, C., Purtscheller, F., Richter, W., and Thoni, M., 1973, Relations between metamorphism and orogeny in a typical section of the Indian Himalayas: *Tschermak's Mineralogische und Petrographische Mitteilungen*, v. 20, p. 303–332, doi: 10.1007/BF01081339.
- Frank, W., Grasemann, B., Guntli, P., and Miller, C., 1995, Geological map of the Kishtwar-Chamba-Kulu region (NW Himalayas, India): *Jahrbuch der Geologischen Bundesanstalt*, v. 138, p. 299–308.
- Grujic, D., Casey, M., Davidson, C., Hollister, L.S., Kundig, R., Pavlis, T., and Schmid, S., 1996, Ductile extrusion of the Higher Himalayan Crystalline in Bhutan: Evidence from quartz microfabrics: *Tectonophysics*, v. 260, p. 21–43, doi: 10.1016/0040-1951(96)00074-1.
- Grujic, D., Hollister, L.S., and Parrish, R.R., 2002, Himalayan metamorphic sequence as an orogenic channel: Insight from Bhutan: *Earth and Planetary Science Letters*, v. 198, p. 177–191, doi: 10.1016/S0012-821X(02)00482-X.
- Heim, A., and Gansser, A., 1939, Central Himalaya Geological Observations of the Swiss Expedition 1936: Zurich, Gebrüder Fretz, 246 p.
- Hodges, K.V., Parrish, R.R., and Searle, M.P., 1996, Tectonic evolution of the central Annapurna Range: *Nepalese Himalayas: Tectonics*, v. 15, p. 1264–1291, doi: 10.1029/96TC01791.
- Jain, A.K., Manickavasagam, R.M., and Singh, S., 1999, Collision tectonics in the NW Himalaya: Deformation, metamorphism, emplacement of leucogranite along Beas-Parbati Valleys, Himachal Pradesh: *Gondwana Research: Group Memoir*, v. 6, p. 3–37.
- Le Fort, P., 1975, Himalayas: The collided range. Present knowledge of the continental arc: *American Journal of Science*, v. 275-A, p. 1–44.
- Le Fort, P., 1996, Evolution of the Himalaya, in Yin, A., and Harrison, T.M., eds., *The Tectonic Evolution of Asia*: New York, Cambridge University Press, p. 95–106.
- Nelson, K.D., and 27 others, 1996, Partially molten middle crust beneath southern Tibet: Synthesis of Project INDEPTH results: *Science*, v. 274, p. 1684–1688, doi: 10.1126/science.274.5293.1684.
- Patel, R.C., Singh, S., Asokan, A., Manickavasagam, R.M., and Jain, A.K., 1993, Extensional tectonics in the Himalayan orogen, Zaskar, NW India, in Treloar, P.J., and Searle, M.P., eds., *Himalayan Tectonics: Geological Society of London Special Publication 74*, p. 445–459.
- Pogue, K.R., Hylland, M.D., Yeats, R.S., Khattak, W.U., and Hussain, A., 1999, Stratigraphic and structural framework of Himalayan foothills, northern Pakistan, in Macfarlane, A., Sorkhabi, R.B., and Quade, J., eds., *Himalaya and Tibet: Mountain Roots to Mountain Tops: Geological Society of America Special Paper 328*, p. 257–274.
- Price, R.A., 1986, The southeastern Canadian Cordillera: Thrust faulting, tectonic wedging, and delamination of the lithosphere: *Journal of Structural Geology*, v. 8, p. 239–254.
- Robinson, D.M., DeCelles, P.G., and Copeland, P., 2006, Tectonic evolution of the Himalayan thrust belt in western Nepal: Implications for channel flow models: *Geological Society of America Bulletin*, v. 118, p. 865–885, doi: 10.1130/B25911.1.
- Searle, M.P., Waters, D.J., Dransfield, M.W., Stephenson, B.J., Walker, C.B., Walker, J.D., and Rex, D.C., 1999, Thermal and mechanical models for the structural and metamorphic evolution of the Zaskar High Himalaya, in Mac Niocall, C., and Ryan, P.D., eds., *Continental Tectonics: Geological Society of London Special Publication 164*, p. 139–156.
- Sharma, V.P., 1977, *Geology of the Kulu-Rampur belt, Himachal Pradesh: Memoirs of the Geological Survey of India*, v. 106, p. 235–407.
- Steck, A., 2003, *Geology of the NW Indian Himalaya: Eclogae Geologicae Helvetiae*, v. 96, p. 147–196.
- Thakur, V.C., 1998, Structure of the Chamba Nappe and position of the main central thrust in Kashmir Himalaya: *Journal of Asian Earth Sciences*, v. 16, p. 269–282, doi: 10.1016/S0743-9547(98)00011-7.
- Vaidya, K.S., 1980, *Geology of Kumaun Lesser Himalaya: Dehra Dun, India, Wadia Institute of Himalayan Geology*, 291 p.
- Vannay, J.C., and Grasemann, B., 1998, Inverted metamorphism in the High Himalaya of Himachal Pradesh (NW India): Phase equilibria versus thermobarometry: *Schweizerische Mineralogische und Petrographische Mitteilungen*, v. 78, p. 107–132.
- Vannay, J.C., and Grasemann, B., 2001, Himalayan inverted metamorphism and syn-convergence extension as a consequence of a general shear extrusion: *Geological Magazine*, v. 138, p. 253–276, doi: 10.1017/S0016756801005313.
- Vannay, J.C., and Steck, A., 1995, Tectonic evolution of the High Himalaya in Upper Lahul (NW Himalaya, India): *Tectonics*, v. 14, p. 253–263, doi: 10.1029/94TC02455.
- Vannay, J.C., Grasemann, B., Rahn, M., Frank, W., Carter, A., Baudraz, V., and Cosca, M., 2004, Miocene to Holocene exhumation of metamorphic crustal wedges in the NW Himalaya: Evidence for tectonic extrusion coupled to fluvial erosion: *Tectonics*, v. 23, p. TC1014, doi: 10.1029/2002TC001429.
- Wyss, M., 2000, Metamorphic evolution of the northern Himachal Himalaya: Phase equilibria constraints and thermobarometry: *Schweizerische Mineralogische und Petrographische Mitteilungen*, v. 80, p. 317–350.
- Wyss, M., Hermann, J., and Steck, A., 1999, Structural and metamorphic evolution of the northern Himachal Himalaya, NW India (Spiti-eastern Lahul-Parvati valley traverse): *Eclogae Geologicae Helvetiae*, v. 92, p. 3–44.
- Yeats, R.S., and Lawrence, R.D., 1984, Tectonics of the Himalayan thrust belt in northwest Pakistan, in Haq, B.U., and Milliman, J.D., eds., *Marine Geology and Oceanography of Arabian Sea and Coastal Pakistan: New York, Van Nostrand Reinhold*, p. 177–198.
- Yin, A., 2006, Cenozoic tectonic evolution of the Himalayan orogen as constrained by along-strike variation of structural geometry, exhumation history, and foreland sedimentation: *Earth-Science Reviews*, v. 76, p. 1–131, doi: 10.1016/j.earsciev.2005.05.004.
- Yin, A., Harrison, T.M., Ryerson, F.J., Chen, W., Kidd, W.S.F., and Copeland, P., 1994, Tertiary structural evolution of the Gangdese thrust system, southeastern Tibet: *Journal of Geophysical Research*, v. 99, p. 18,175–18,201, doi: 10.1029/94JB00504.

Manuscript received 29 March 2007

Revised manuscript received 4 June 2007

Manuscript accepted 8 June 2007

Printed in USA